

Complete Genome Sequence of Staphylococcus aureus Siphophage  
Lorac

Antoine Marc et al.

Deposited 2023-09-27

Citation of published version:

Marc, A., Cater, K., Kongari, R., Hatoum-Aslan, A., Young, R. F., III, & Liu, M.  
(2019). Complete Genome Sequence of Staphylococcus aureus Siphophage Lorac.  
In J. J. Dennehy (Ed.), *Microbiology Resource Announcements* (Vol. 8, Issue 27).  
American Society for Microbiology. <https://doi.org/10.1128/mra.00586-19>

©2019

This work is licensed under a Attribution 4.0 International (CC BY 4.0) license.





# Complete Genome Sequence of *Staphylococcus aureus* Siphophage Lorac

Antoine Marc,<sup>a</sup> Katie Cater,<sup>b</sup>  Rohit Kongari,<sup>a</sup> Asma Hatoum-Aslan,<sup>b</sup> Ryland F. Young III,<sup>a</sup> Mei Liu<sup>a</sup>

<sup>a</sup>Center for Phage Technology, Texas A&M University, College Station, Texas, USA

<sup>b</sup>Department of Biological Sciences, The University of Alabama, Tuscaloosa, Alabama, USA

**ABSTRACT** *Staphylococcus aureus* is a leading cause of a wide range of clinical infections. Here, we announce the complete genome sequence of *S. aureus* siphophage Lorac, a phiETA-like temperate phage that is similar at the nucleotide level to the previously described *S. aureus* prophage phiNM2.

*Staphylococcus aureus* is a Gram-positive skin-associated bacterium and is the causative agent for a wide range of clinical infections, including bacteremia, infective endocarditis, and pleuropulmonary and other device-related infections (1). Temperate phages, such as phiETA, are able to move between *S. aureus* strains and can encode virulence factors or toxins (2).

Siphophage Lorac was isolated from the wastewater treatment plant in Tuscaloosa, Alabama, in August 2015 using the host strain *S. aureus* strain RN4220. Host bacteria were cultured on tryptic soy broth or agar (Difco) at 37°C with aeration. Phages were cultured and propagated by the soft agar overlay method (3). It was identified as a siphophage using negative-stain transmission electron microscopy performed at the University of Alabama Optical Analysis Facility, as described previously (4). Phage genomic DNA was prepared using a modified Promega Wizard DNA cleanup kit protocol as described previously (4). Pooled indexed DNA libraries were prepared using the Illumina TruSeq Nano low-throughput (LT) kit, and the sequence was obtained from the Illumina MiSeq platform using the MiSeq v2 500-cycle reagent kit following the manufacturer's instructions, producing 631,646 reads for the index containing the phage genome. FastQC 0.11.5 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) was used to quality control reads. The reads were trimmed with FastX-Toolkit 0.0.14 ([http://hannonlab.cshl.edu/fastx\\_toolkit/download.html](http://hannonlab.cshl.edu/fastx_toolkit/download.html)) before being assembled using SPAdes 3.5.0 (5). Contig completion was confirmed by PCR using primers (5'-GTCCCTATCAAACCGAGAATCC-3' and 5'-ACATGGGTGTAATCGACAAAGA-3') facing off the ends of the assembled contig and Sanger sequencing of the resulting product, with the contig sequence manually corrected to match the resulting Sanger sequencing read. GLIMMER 3.0 (6) and MetaGeneAnnotator 1.0 (7) were used to predict protein-coding genes with manual correction for appropriate gene starts, and tRNA genes were predicted with ARAGORN 2.36 (8). Rho-independent termination sites were identified via TransTerm (<http://transterm.cbcb.umd.edu/>). Sequence similarity searches were performed by BLASTp 2.2.28 (9) against the NCBI nonredundant (nr), UniProt Swiss-Prot (10), and TrEMBL databases. InterProScan 5.15-54.0 (11), LipoP (12), and TMHMM v2.0 (13) were used to predict protein function. All analyses were conducted at default settings via the CPT Galaxy (14) and WebApollo (15) interfaces (<https://cpt.tamu.edu/galaxy-pub>).

Lorac was assembled as a complete genome with 43,147 bp and 1,371-fold coverage. It has a G+C content of 34%, which is similar to that of its host (16). Lorac is probably a temperate phage since it belongs to the *Phietavirus* genus and is 99.18%

**Citation** Marc A, Cater K, Kongari R, Hatoum-Aslan A, Young RF, III, Liu M. 2019. Complete genome sequence of *Staphylococcus aureus* siphophage Lorac. *Microbiol Resour Anounc* 8:e00586-19. <https://doi.org/10.1128/MRA.00586-19>.

**Editor** John J. Dennehy, Queens College

**Copyright** © 2019 Marc et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Asma Hatoum-Aslan, [asma.hatoum@ua.edu](mailto:asma.hatoum@ua.edu), or Mei Liu, [meiliu@tamu.edu](mailto:meiliu@tamu.edu).

**Received** 17 May 2019

**Accepted** 30 May 2019

**Published** 3 July 2019

similar at the nucleotide level to phiNM2 (GenBank accession no. [DQ530360](https://doi.org/10.1093/nar/33.11.2105)), a prophage identified in *S. aureus* strain Newman (17). Morphogenesis genes coding for the major capsid protein, scaffolding protein, TerL, TerS, tail proteins, tail-head connector protein, portal protein, tape measure protein, and tape measure chaperone were annotated. Lysogeny-associated proteins, including Cro-like and cI-like regulators, excisionase, and integrase, were identified, as was a lysis cassette consisting of a class II holin and an amidase-type endolysin.

**Data availability.** The genome sequence of phage Lorac was submitted to GenBank under accession no. [MH321492](https://doi.org/10.1093/nar/42.11.2105). Associated BioProject, SRA, and BioSample accession numbers are [PRJNA222858](https://doi.org/10.1093/nar/42.11.2105), [SRR8788599](https://doi.org/10.1093/nar/42.11.2105), and [SAMN11260821](https://doi.org/10.1093/nar/42.11.2105), respectively.

## ACKNOWLEDGMENTS

This work was supported by funding from the National Science Foundation (awards EF-0949351, DBI-1565146, and MCB-1749886). Additional support came from the Center for Phage Technology (CPT), an Initial University Multidisciplinary Research Initiative supported by Texas A&M University and Texas AgriLife, and from the Department of Biochemistry and Biophysics at Texas A&M University.

We are grateful for the advice and support of the CPT staff and Kim Lackey at the University of Alabama Optical Analysis Facility.

This phage was discovered in a pilot run for BSC 411 Phage Discovery, an undergraduate course at the University of Alabama. This announcement was prepared in partial fulfillment of the requirements for BICH464 Phage Genomics, an undergraduate course at Texas A&M University.

## REFERENCES

- Tong SYC, Davis JS, Eichenberger E, Holland TL, Fowler VG, Jr. 2015. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 28:603–661. <https://doi.org/10.1128/CMR.00134-14>.
- Yamaguchi T, Hayashi T, Takami H, Nakasone K, Ohnishi M, Nakayama K, Yamada S, Komatsuzawa H, Sugai M. 2000. Phage conversion of exfoliative toxin A production in *Staphylococcus aureus*. *Mol Microbiol* 38:694–705. <https://doi.org/10.1046/j.1365-2958.2000.02169.x>.
- Adams MK. 1959. Bacteriophages. Interscience Publishers Inc., New York, NY.
- Cater K, Dandu VS, Bari SM, Lackey K, Everett GF, Hatoum-Aslan A. 2017. A novel *Staphylococcus podophage* encodes a unique lysin with unusual modular design. *mSphere* 2:e00040-17. <https://doi.org/10.1128/mSphere.00040-17>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27:4636–4641. <https://doi.org/10.1093/nar/27.23.4636>.
- Noguchi H, Taniguchi T, Itoh T. 2008. MetaGeneAnnotator: detecting species-specific patterns of ribosomal binding site for precise gene prediction in anonymous prokaryotic and phage genomes. *DNA Res* 15:387–396. <https://doi.org/10.1093/dnares/dsn027>.
- Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
- The UniProt Consortium. 2018. UniProt: the universal protein knowledgebase. *Nucleic Acids Res* 46:2699. <https://doi.org/10.1093/nar/gky092>.
- Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30:1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>.
- Juncker AS, Willenbrock H, Von Heijne G, Brunak S, Nielsen H, Krogh A. 2003. Prediction of lipoprotein signal peptides in Gram-negative bacteria. *Protein Sci* 12:1652–1662. <https://doi.org/10.1110/ps.0303703>.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 305:567–580. <https://doi.org/10.1006/jmbi.2000.4315>.
- Cock PJ, Gruning BA, Paszkiewicz K, Pritchard L. 2013. Galaxy tools and workflows for sequence analysis with applications in molecular plant pathology. *PeerJ* 1:e167. <https://doi.org/10.7717/peerj.167>.
- Lee E, Helt GA, Reese JT, Munoz-Torres MC, Childers CP, Buels RM, Stein L, Holmes IH, Elisk CG, Lewis SE. 2013. Web Apollo: a Web-based genomic annotation editing platform. *Genome Biol* 14:R93. <https://doi.org/10.1186/gb-2013-14-8-r93>.
- Suzuki H, Lefebure T, Bitar PP, Stanhope MJ. 2012. Comparative genomic analysis of the genus *Staphylococcus* including *Staphylococcus aureus* and its newly described sister species *Staphylococcus simiae*. *BMC Genomics* 13:38. <https://doi.org/10.1186/1471-2164-13-38>.
- Bae T, Baba T, Hiramatsu K, Schneewind O. 2006. Prophages of *Staphylococcus aureus* Newman and their contribution to virulence. *Mol Microbiol* 62:1035–1047. <https://doi.org/10.1111/j.1365-2958.2006.05441.x>.